

Astronomical Journal, in press (January 2003)

**Addendum: “Hubble Space Telescope Evidence for an Intermediate-Mass
Black Hole in the Globular Cluster M15—
II. Kinematical Analysis and Dynamical Modeling¹”**

Joris Gerssen, Roeland P. van der Marel

Space Telescope Science Institute, 3700 San Martin Drive, Baltimore, MD 21218

Karl Gebhardt

Department of Astronomy, Mail Code C4100, University of Texas at Austin, Austin, TX 78712

Puragra Guhathakurta^{2,3}

Herzberg Institute of Astrophysics, National Research Council of Canada, 5071 West Saanich
Road, Victoria, BC V9E 2E7, Canada

Ruth C. Peterson⁴

UCO/Lick Observatory, Department of Astronomy and Astrophysics, University of California at
Santa Cruz, 1156 High Street, Santa Cruz, CA 95064

Carlton Pryor

Department of Physics and Astronomy, Rutgers University, 136 Frelinghuysen Road, Piscataway,
NJ 08854-8019

¹Based on observations made with the NASA/ESA Hubble Space Telescope, obtained at the Space Telescope Science Institute, which is operated by the Association of Universities for Research in Astronomy, Inc., under NASA contract NAS 5-26555. These observations are associated with proposal #8262.

²Herzberg Fellow.

³Permanent address: UCO/Lick Observatory, University of California, Santa Cruz, 1156 High Street Santa Cruz, California 95064.

⁴Also at: Astrophysical Advances, Palo Alto, CA 94301.

In our paper we reported the existence of a dark and compact mass component near the center of M15, based on an analysis of new data from the Hubble Space Telescope. Possible explanations for this mass component include: (1) a single intermediate-mass black hole (BH); or (2) a collection of dark remnants (e.g., neutron stars) that have sunk to the cluster center due to mass segregation. We assessed the plausibility of the latter possibility by comparing the kinematical data for M15 to the predictions of the most sophisticated and most recently published Fokker-Planck models for M15 (Dull et al. 1997, hereafter D97). We showed that the mass-to-light ratio (M/L) profile in Figure 12 of D97 implies too few dark remnants near the cluster center to explain the observed kinematics of M15. This supported the view that M15 harbors an intermediate-mass BH. We address here how this conclusion is affected by the recent discovery of an error in Figure 12 of D97. We show that the presence of an intermediate-mass BH continues to be a viable interpretation of the data, but that its presence ceases to be uniquely implied.

After the completion of our paper, the authors of the D97 paper discovered an unfortunate error in their Figures 9 and 12. The labeling along the abscissa of these figures is incorrect due to a coding error in SM plotting routines (H. Cohn and B. Murphy, private communication, 2002). The units along the top axis should have read ‘arcmin’ instead of ‘pc’, and the labeling in arcmin along the bottom axis is incorrect. The net result is that the radial scale of these figures in the D97 paper is too compressed by a factor 2.82. The true total mass of the centrally concentrated population of dark remnants in the Fokker-Planck models is therefore considerably larger than what was implied by the M/L profile shown in Figure 12 of the D97 paper.

Figure 1 shown here is similar to Figure 12 of our paper, but it now shows the data-model comparison with the corrected D97 M/L profile. Models without a BH are now found to be statistically acceptable (within 1σ), although inclusion of an intermediate-mass BH, with $M_{\text{BH}} = 1.7^{+2.7}_{-1.7} \times 10^3 M_{\odot}$, still provides a marginally better fit to the data (although not a statistically significant one). Hence, the Fokker-Planck models for M15 discussed by D97 do in fact have enough dark remnants near the cluster center to explain the observed kinematics. However, the D97 models still have a number of important shortcomings, as discussed in §5.4 of our paper. Most importantly, D97 assumed that all neutron stars that form in the cluster are retained. By contrast, the observed distribution of pulsar kick velocities indicates that the retention factor should only be a few percent; most authors agree that it should be no more than 10% (see references in §5.4 of our paper). In this sense, the D97 models provide an upper limit on the number and mass of dark remnants in M15. The same is true for N -body models constructed recently by Baumgardt et al. (2002), the results of which are qualitatively similar to those of D97. More realistic evolutionary models that include neutron star escape will require a more massive and more statistically significant BH to fit the data than that suggested by Figure 1. Alternatively, one can assume that there are more stars in the high-mass end of the initial mass function, or that the transition between stars that evolve to white dwarfs as compared to neutron stars occurs at a higher initial mass (H. Cohn and B. Murphy, private communication, 2002). Independent evidence does not exist to support these assumptions, although they cannot be ruled out.

The evidence for a central BH in M15 is less convincing than it was on the basis of our earlier analyses. However, because of the somewhat unrealistic assumptions about neutron star retention in the models of D97 and Baumgardt et al. (2002), and because of the independent evidence for a

BH in another cluster (G1; Gebhardt, Rich & Ho 2002), the presence of a BH in M15 continues to be a viable interpretation of the data. The best fit BH mass with the corrected D97 M/L profile is $M_{\text{BH}} = 1.7^{+2.7}_{-1.7} \times 10^3 M_{\odot}$ (see Figure 1); with a constant M/L it is $M_{\text{BH}} = 3.2^{+2.2}_{-2.2} \times 10^3 M_{\odot}$ (see §5.4 of our paper). A model that includes both neutron star escape and mass segregation would probably yield a value between these numbers. So if M15 has a BH, its mass is consistent with the correlation between velocity dispersion and BH mass that has been inferred for galaxies (see Figure 14 of our paper). This continues to suggest the possible existence of an important new link between the structure, evolution and formation of globular clusters, galaxies, and their central BHs (see §6 of our paper, and also Gebhardt et al. 2002). However, with the presently available models and data it is neither uniquely implied nor ruled out that M15 has an intermediate-mass BH.

We thank Haldan Cohn, Brian Murphy, Phyllis Lugger, Piet Hut and Steve McMillan for stimulating discussions.

REFERENCES

- Baumgardt, H., Hut, P., Makino, J., McMillan, S., & Portegies Zwart, S. 2002, ApJL, submitted
- Dull, J. D., Cohn, H. N., Lugger, P. M., Murphy, B. W., Seitzer, P. O., Callanan, P. J., Rutten, R. G. M., & Charles, P. A. 1997, ApJ, 481, 267 (D97)
- Gebhardt, K., Rich, R. M., & Ho, L. 2002, ApJ, 578, L41




Fig. 1.— Data-model comparison for spherical dynamical models with an isotropic velocity distribution, an M/L profile inferred from Fokker-Planck models, and a central black hole of mass M_{BH} . The M/L profile is a corrected version of the one shown by D97. **(a; left panel)** The likelihood quantity λ defined in equation (4) of our paper as function of M_{BH} . The minimum in λ identifies the best fit black hole mass. Horizontal dashed lines indicate the 1 and 2σ confidence regions. **(b; right panel)** The RMS projected line-of-sight velocity σ_{RMS} as a function of projected radius R . The heavy jagged curve surrounded by heavy dashed curves is the observed profile, as in Figure 9d of our paper. The smooth thin curves are the predictions for models with M_{BH} ranging from 0 to $10 \times 10^3 M_{\odot}$ in steps of $10^3 M_{\odot}$.